

APPROVED: SIGNATURE ON FILE IN EMS OFFICE
Executive Director

SIGNATURE ON FILE IN EMS OFFICE
Medical Director

EFFECTIVE DATE: 01/01/2004

SUPERSEDES: _____

REVISED: _____

REVIEW DATE: 01/01/2009

PAGE: 1 OF 1

PEDIATRIC DYSTONIC REACTION TO PHENOTHIAZINE DRUGS

- I. AUTHORITY: Health and Safety Code, Division 2.5, California Code of Regulations, Title 22, Division 9
- II. PURPOSE: To serve as the treatment standard for EMT-Is and EMT-Ps in treating patients.
- III. PROTOCOL:

History of ingestion of phenothiazine with restlessness (akathesias), muscle spasms of the neck, jaw and back, movement of eyeballs (oculogyric crisis), frightened, small pupils, facial grimace, protruding tongue, opisthotonus (back arching).

Phenothiazines are prescribed for their antiemetic and tranquilizing properties. Phenothiazines include: chlorpromazine (Thorazine), metoclopramide (Reglan), prochlorperazine (Compazine), promethazine (Phenergan and Atarax).

Other medications that can cause dystonic reactions include: droperidol (Inapsine) and haloperidol (Haldol).

NOTE: Phenothiazine reactions may occur at normal dosing levels and administration of charcoal is not necessary.

STANDING ORDERS

ABC's

OXYGEN

MONITOR

Treat rhythm as appropriate.

IV/IO ACCESS

TKO with microdrip tubing and volume control chamber.

DIPHENHYDRAMINE

1 mg/kg IV push (maximum dose 50 mg) or IM if IV access not promptly available. May repeat if needed.