

APPROVED: Signature On File In EMS Office
Executive Director

Signature On File In EMS Office
Medical Director

EFFECTIVE DATE: 7/01/2011
SUPERSEDES: _____
REVISED: _____
REVIEW DATE: 7//2016
PAGE: 1 OF 1

PEDIATRIC SYMPTOMATIC BRADYCARDIA

- I. AUTHORITY: Health and Safety Code, Division 2.5, California Code of Regulations, Title 22, Division 9
- II. PURPOSE: To serve as the treatment standard for EMTs and Paramedics in treating patients.
- III. PROTOCOL: Most bradycardias in children are due to hypoxia. Bradycardia may also be secondary to sinus node disease, increased parasympathetic tone or drug effects (e.g., digitalis, beta-blockers, or calcium antagonists), hypothermia or head injury. Heart rate is below 60 beats per minute, with associated signs/symptoms of low cardiac output. Never treat any bradycardia if the patient does not have serious symptoms.

STANDING ORDERS

ASSESS	CAB
OXYGEN	NOTE: Most bradycardias in children are due to hypoxia.
SECURE AIRWAY	Using the simplest effective method. A BLS airway with objective evidence of good ventilation and oxygenation is adequate and acceptable. Consider intubation only if unable to establish adequate ventilation and oxygenation using a BVM. Refer to General Procedures Protocol 554.00
MONITOR	
IV/IO ACCESS	TKO with microdrip tubing and volume control chamber.
ASSESS	For: 1. Heart rate: less than 80 beats per minute in infants (less than 1 year); less than 60 beats per minute in children (1 - 12 years). AND 2. Signs of poor perfusion (delayed capillary refill, diminished distal pulses, cool extremities, altered level of consciousness) or respiratory distress.
CPR	If heart rate less than 80/minute in infant or less than 60/minute child.
EPINEPHRINE	0.01 mg/kg of 1:10,000 IV/IO. Repeat every 3 minutes until above heart rate target or signs of poor perfusion or respiratory distress have improved.
CONSIDER	
ATROPINE	0.02 mg/kg IV/IO. Minimum dose 0.1 mg. Maximum single dose 0.5 mg. May be repeated once in 3 minutes.